

- ¹⁵ College of Physicians and Surgeons of Ontario. (2000). *Evidence-Based Recommendations for Medical Management of Chronic Non-Malignant Pain*. Retrieved from http://www.cpsa.ab.ca/Libraries/pro_tpp/MedicalManagementPain.pdf?sfvrsn=0. Accessed September 24, 2013.
- ¹⁶ World Health Organization. (2013). *WHO's Pain Ladder for Adults*. Retrieved from <http://www.who.int/cancer/palliative/painladder/en/>. Accessed October 10, 2013.
- ¹⁷ Federation of State Medical Boards of the United States, Inc. (2004). *Model Policy for the Use of Controlled Substances for the Treatment of Pain*. Retrieved from http://www.fsmb.org/pdf/2004_grpol_controlled_substances.pdf. Accessed September 24, 2013.
- ¹⁸ Berndt, S., Maier, C., & Schutz, H.W. (1993). Polymedication and medication compliance in patients with chronic non-malignant pain. *Pain*, 52(3), 331-339.
- ¹⁹ Trescot, A.M., Helm, S., Hansen, H., Benyamin, R., Glaser, S.E., Adlaka, R., . . . Manchikanti, L. (2008). Opioids in the management of chronic non-cancer pain: An update of American society of the interventional pain physicians' (ASIPP) guidelines. *Pain Physician*, 11, S5-S62.
- ²⁰ Daniell, H.W. (2008). Opioid endocrinopathy in women consuming prescribed sustained-action opioids for control of nonmalignant pain. *Journal of Pain*, 9 (1), 28-36.
- ²¹ Passik, S.D. & Weinreb, H.J. (2000). Managing chronic nonmalignant pain: Overcoming obstacles to the use of opioids. *Advances in Therapy*, 17(2), 70-83.
- ²² Quang-Cantagrel, N.D., Wallace, M.S., & Magnuson, S.K. (2000). Opioid substitution to improve the effectiveness of chronic noncancer pain control: A chart review. *Anesthesia and Analgesia*, 90(4), 933-937.
- ²³ Smith, H.S. (2009). *Current therapy in pain*. Philadelphia, PA: Saunders Elsevier.
- ²⁴ CredibleMeds powered by AZCERT.org. (2013). *Composite List of All QT Drugs and the List of Drugs to Avoid for Patients with Congenital LQTS*. Retrieved from <http://www.crediblemeds.org/everyone/composite-list-all-qtdrugs/>. Accessed September 24, 2013.
- ²⁵ Walker, J.M., Farney, R.J., Rhondeau, S.M., Boyle, K.M., Valentine, K., Cloward, T.V., & Shilling, K.C. (2007). Chronic opioid use is a risk factor for the development of central sleep apnea and ataxic breathing. *Journal of Clinical Sleep Medicine*, 3(5), 455-461.