

	HEPATITIS A	HEPATITIS B	HEPATITIS C
<b>MODES OF TRANSMISSION</b>	<ul style="list-style-type: none"> <li>• Close person-to-person contact with an infected person</li> <li>• Sexual contact with an infected person</li> <li>• Ingestion of contaminated food or water</li> </ul>	<ul style="list-style-type: none"> <li>• Blood and other body fluids containing HBV</li> <li>• Sharing IDU equipment</li> <li>• Perinatal transmission</li> <li>• Unprotected sex</li> <li>• Accidental needle sticks</li> <li>• Blood transfusion (rare)</li> </ul>	<ul style="list-style-type: none"> <li>• Sharing IDU equipment</li> <li>• Unprotected sex</li> <li>• Perinatal transmission</li> </ul>
<b>INCUBATION PERIOD</b>	15-50 days (average 28 days)	60-150 days (average 90 days)	14-182 days (average 14-84 days)
<b>% OF NEWLY INFECTED ADULTS WHO DEVELOP CHRONIC INFECTION</b>	0%	2-6%	75-85%
<b>IMMUNITY AFTER INFECTION</b>	Yes	Yes	No
<b>VACCINE AVAILABLE</b>	Yes	Yes	No
<b>TREATMENT</b>	Supportive care	Yes, but not curative	Yes, curative