


Oklahoma Geological Survey Guidebook 22



Guide to
**Robbers Cave
State Park**

Arthur J. Myers, Dearl T. Russell
George J. Goodman, Cheryl A. Lawson



Oklahoma Geological Survey
Charles J. Mankin, *Director*

ISSN 0078-4400

Guidebook 22

GUIDE TO ROBBERS CAVE STATE PARK

**Arthur J. Myers, Dearl T. Russell
George J. Goodman, Cheryl A. Lawson**

**The University of Oklahoma
Norman**

1986

FOREWORD

An earlier edition of *Guide to Robbers Cave State Park* was published by the Oklahoma Geological Survey (OGS) in 1958 as Guidebook 7. It contained information on the geology and biology of Robbers Cave State Park and the surrounding area. The author of Guidebook 7, Dearl T. Russell, had studied and mapped northern Latimer County and submitted a thesis on his investigations for a Master of Science degree at the University of Oklahoma in 1958. He prepared Guidebook 7 to help Scouts at Camp Tom Hale and visitors to Robbers Cave enjoy and understand the natural history of the area.

The original material has been updated and expanded by Dr. Arthur J. Myers, Dr. George J. Goodman, and Dr. Cheryl A. Lawson, and is now being issued as Guidebook 22 by the OGS.

Myers retired from the OGS in 1984 after serving six years as a geomorphologist and aerial-photo interpreter. Before coming to the OGS, he had served on the faculty of the University of Oklahoma School of Geology and Geophysics for 33 years. Myers holds a B.A. in chemistry from Kalamazoo College, a B.S. and M.S. in geological engineering from Michigan Technological University, Houghton, and a Ph.D. in geology from the University of Michigan.

The botany section of this publication was prepared by Goodman and Lawson. Goodman, a recognized authority on the flora of Oklahoma, is a Regents Professor Emeritus of Botany at the University of Oklahoma and was formerly curator of the Bebb Herbarium there for more than 30 years. His current research is on the flora of Oklahoma and the botany and botanical history of the western United States.

Lawson graduated with distinction from the University of Oklahoma and from there received her Ph.D. in botany. She is a member of Phi Beta Kappa, and of many state, national, and international professional organizations. Her current research involves the botany of the Great Plains and the Rocky Mountains.

Cover Illustration

Robbers Cave

Leaf drawings by Roy D. Davis

This publication, printed by the University of Oklahoma Printing Services, Norman, Oklahoma, is issued by the Oklahoma Geological Survey as authorized by Title 70, Oklahoma Statutes, 1981, Section 3310, and Title 74, Oklahoma Statutes, 1981, Sections 231–238. 1,500 copies have been prepared for distribution at a cost of \$5,678.00 to the taxpayers of the State of Oklahoma. Copies have been deposited with the Publications Clearinghouse of the Oklahoma Department of Libraries.

CONTENTS

	<i>Page</i>
Geology of Robbers Cave State Park	1
Introduction	1
Origin of Caves	1
Origin of Rocks	5
Geologic History	6
Stratigraphy	7
Paleontology	8
Topography and Structural Geology	10
Botany of Robbers Cave State Park	19
Introduction	19
Key to Woody Plants	21
List of Plants	36
References	46
Glossary of Geological Terms	47

FIGURES

1. Map of Robbers Cave State Park area and nearby highways in southeastern Oklahoma	2
2. Location map of available facilities at Robbers Cave State Park	3
3. Photograph of Robbers Cave	4
4. Geologic time scale compared to calendar year	7
5. Geologic provinces of Oklahoma	8
6. Columnar section, Robbers Cave area	9
7. <i>Stigmaria</i>	10
8. <i>Calamites</i>	10
9. Scale trees	11
10. Geomorphic regions of Oklahoma and parts of surrounding states	13
11. Geomorphic provinces in central and eastern Oklahoma	15
12. Aerial view of canoe-shaped mountain west of Robbers Cave State Park	15
13a. Topographic map of Tucker Knob	16
13b. Tucker Knob	16
14. Bluff at Robbers Cave	17
15. Joints in sandstone	17
16. Slumped blocks of sandstone	17
17. Small cuesta in the Robbers Cave area	18
18. Lake Carlton	18
19. Illustrations showing leaf organization	25
20. Illustrations showing leaf parts, position, and margins	26
21. Illustrations showing simple, alternate leaved species	27
22. Illustrations showing simple, alternate leaved species	28
23. Illustrations showing simple, alternate leaved species	29
24. Illustrations showing simple, alternate leaved species	30
25. Illustrations showing simple, alternate leaved species	31
26. Illustrations showing compound, alternate leaved species	32
27. Illustrations showing compound, alternate leaved species	33
28. Illustrations showing compound, alternate leaved species	34
29. Illustrations showing simple, opposite leaved species	35

PLATE

1. Geologic map of Robbers Cave State Park area	pocket
---	--------

Geology of Robbers Cave State Park

Arthur J. Myers¹ and Dearl T. Russell²

Introduction

Robbers Cave State Park is in the wooded hills of southeastern Oklahoma. It is on the northern flank of the Ouachita Mountains, where sandstones of Pennsylvanian age, interbedded with shales, form rugged and picturesque landscapes. Figure 1 is a map showing the location of the park in northwestern Latimer County.

The park is a forested area that is refreshing and enjoyable, especially for persons from the grasslands of central and western Oklahoma. It offers recreational facilities of many kinds (fig. 2), including several hiking trails from which to observe trees, rocks, and wildlife. Some of the trails are short, but for energetic, adventuresome people there is a 12-mile hiking trail from Robbers Cave to the picnic area south of Lake Carlton.

Both Wayne Wallace Reservoir and Lake Carlton are formed by dams on Fourche Maline Creek and are picturesque bodies of water with high bluffs (formed by the Bluejacket Sandstone of Pennsylvanian age) on their far sides. As it flows through the park from Wayne Wallace Reservoir to Lake Carlton, Fourche Maline Creek has many interesting and scenic points. The water flowing over the dam at Lake Carlton is pretty, but even more important on a hot summer day, Lake Carlton boasts a swimming beach as well as a swimming pool.

Lodging facilities include cabins along the escarpment side of a cuesta with a lovely view of Coon Creek Valley and Coon Creek Lake. Trailer hookups

are available in the Old Circle Campground area. Group-camp cabins are available as well, and there are three campgrounds where tents can be pitched. The park even has an equestrian campground. Many picnic shelters are available, along with a cafe and grocery store.

Of course, one of the main attractions of the park is Robbers Cave (fig. 3). From the parking area a path leads to the cave, which appears rather small and insignificant at first and may be an initial disappointment to some. But very few visitors leave the park without succumbing to the temptation to climb over the rocks and boulders to explore the cave.

Origin of Caves

Man has long been fascinated by caves. His first shelter was a cave, and the caves at Lascaux, France, contain some of the earliest works of art, the wall paintings that were made by Cro-Magnon man more than 20,000 years ago. The Dead Sea Scrolls reposed for centuries in the silent security of caves, which their ancient custodians recognized as suitable repositories for their preservation and eventual recovery. Caves are often tourist attractions; witness the popularity of Luray Cavern in Virginia, Mammoth Cave in Kentucky, Onondaga and

¹Geologist (retired), Oklahoma Geological Survey.

²Deceased.




Figure 1. Map of southeastern Oklahoma, showing routes of major highways and location of Robbers Cave State Park and nearby State parks.

Meramec Caverns in Missouri, and Carlsbad Caverns in New Mexico.

Not everyone is in complete agreement about the detailed mechanics of cave formation. There is agreement that caves are formed when soluble rock is dissolved by subsurface water; although some scientists believe the dissolution takes place above the water table, some believe it is at the water table, and some believe that it is below the water table.

Most caves form in limestone, which is made up of the mineral calcite, whose chemical composition is calcium carbonate or CaCO_3 . Carbonic acid, a weak acid that forms in water where there is decaying organic matter, reacts with the CaCO_3 to form a soluble bicarbonate. As a result, ground water moving along joints and bedding planes will dissolve the rock and form enlargements which, in time, become caves.

Gypsum is another rock that is soluble in water, and again, water moving along joints and bedding

planes can dissolve the rock and form caves. Alabaster Cavern, in Woodward County, Oklahoma, was formed by this method in gypsum.

Robbers Cave formed in sandstone that is composed of the mineral quartz (SiO_2), which is relatively insoluble. However, the quartz grains have been cemented together by calcium carbonate so ground water can dissolve the calcium carbonate cement, and then the loosened quartz grains can be removed by the circulating ground water. The sandstone bluff (IPsv-11 on geologic map, pl. 1, in pocket) in which Robbers Cave has formed rises about 140 feet above the camp's parking area. The cave is approximately two-thirds of the way up the bluff, where it has formed along the junction of a bedding plane and a fault. This was a zone of weakness, and downward-percolating ground water moving along this zone gradually enlarged the opening to form the cave. As the cave became larger, blocks of rock were loosened by dissolution of the cement-




Figure 2. Map of Robbers Cave State Park, showing location of available facilities.




Figure 3. Robbers Cave: the floor is a bedding plane that dips from right to left; the steep slope on the left side represents a fault plane; the roof is irregular because of weathering and slumping. Erosion probably began along the junction of the bedding plane and the fault plane and was enlarged by weathering, slumping, and erosion.

ing material and subsequently fell to the floor of the cave. Some of these blocks were removed later by erosion.

Weathering of some of the minerals has stained many of the rocks shades of red, brown, and various other colors—making some beautiful abstract patterns. Weathering and erosion along bedding planes also has intensified the stratification. In many places a series of steps has formed along individual strata, making for an easy climb toward the top of the bluff. The top of the bluff is a wooded area with many large boulders and, at the edge of the large slump blocks, a spectacular view of the valley to the southeast.

Origin of Rocks

Rocks are classified as sedimentary (formed from the accumulation of sediments, either above or below sea level), igneous (formed from the cooling and solidification of molten material), or metamorphic (formed by alterations in the solid state of pre-existing rock in response to changes in heat and (or) pressure below the zones of weathering or cementation). In Oklahoma, almost 99 percent of the outcropping rocks are sedimentary in origin; the rest are mainly igneous rocks in the Wichita and Arbuckle Mountains, with a small area of mildly metamorphosed rocks in the Ouachita Mountains southeast of Robbers Cave State Park.

At various times in the geologic past, parts of continents were flooded by shallow inland seas. Rivers flowing into these seas transported sand, silt, and dissolved mineral matter, which waves and currents then spread out in layers that are called strata or beds. Rocks formed in this manner from sediments are sedimentary rocks, and their most characteristic feature is the stratification. The tops and bottoms of strata are bedding planes. As originally deposited, the strata are essentially horizontal, and the first one deposited is at the bottom and is the oldest. Successive overlying strata are younger than the ones below. A unit of rock of the same rock type is called a formation, and within a formation there may be several strata. Formations are generally named for localities where they were first studied and described.

Sedimentary rocks are classified on the basis of texture (size of the mineral grains) and (or) mineral composition and are either clastic or nonclastic rocks. Clastic rocks consist of fragments derived from preexisting rocks that were broken down by

weathering and subsequently transported and deposited in another environment. A sandstone is a sedimentary deposit composed of sand-sized (1/16 to 2 mm) quartz grains that have been lithified (made into a rock) by cementation of a secondary mineral (commonly calcite, chemical composition calcium carbonate or CaCO_3). Siltstone is a cemented sedimentary deposit composed of silt-sized (1/256 to 1/16 mm) quartz grains, and it generally contains impurities of clay minerals. Shale is a laminated sedimentary deposit containing clay-sized particles (less than 1/256 mm) that have been lithified, or changed to rock, most commonly by compaction. The sandstones and shales of Latimer County are primarily buff and brown colored because the mineral grains are coated with a thin covering of iron oxide; however, some rocks are gray because of a coating of organic matter.

Nonclastic sedimentary rocks form when dissolved mineral matter is precipitated on the sea floor. Limestone, composed of the mineral calcite, is the most abundant nonclastic sedimentary rock. Limestone is still being formed—in places such as in the seas around the Bahama Islands—by both physiochemical precipitation (direct precipitation from sea water) and biochemical precipitation (removal by marine organisms of dissolved mineral matter from sea water). Limestone formed in this manner can later be broken down by weathering and (or) erosion, then transported by rivers back to the sea, where it is redeposited by waves and currents. Since the material has been broken up and transported from its place of origin, it becomes a clastic limestone. When the loss of water by evaporation from the sea is greater than the amount of water brought to it by rivers, the dissolved mineral matter is precipitated on the sea floor. These deposits are called evaporites, and include gypsum and salt (the mineral halite). Although there are no evaporites in Latimer County, they are abundant in western Oklahoma. In fact, Alabaster Caverns State Park in Woodward County has a cave that is formed in gypsum. In Latimer County, only one thin limestone is in the Robbers Cave area, although a thick limestone bed forms a ridge south of Wilburton.

Coal is another type of sedimentary rock in Latimer County. It consists of compressed carbonaceous matter derived from the partial decomposition of plant life. Plant remains accumulated in swampy waters and were partially preserved in the stagnant waters of large fresh-water swamps. Subsequent burial by more sediments resulted in increased pressure and a rise in temperature that caused the organic plant remains to be transformed into bituminous coal. Coal beds in the park area are thin and low in

grade (level of purity); however, a high-grade bituminous coal is mined from surface pits near Wilburton.

Geologic History

Geology is centered around the history of the Earth and the study of the processes that produced the geologic features seen today. Less than 200 years ago many respected scientists believed that the Earth was only a few thousand years old. James Usher, an Irish archbishop, used genealogies in the Bible as a basis for publishing in "Anneles Veteris et Novi Testamenti" that the Earth was created in 4004 B.C. The vice-chancellor of Cambridge University refined the calculations and determined that the Earth was created at 9:00 a.m. on October 26, 4004 B.C. However, Charles Darwin (1809–1882), after his observations on the voyage of the Royal Navy's H.M.S. Beagle and with the publication of *The Origin of Species*, believed that the Earth had to be much older in order to account for the changes that had taken place. Today it is believed that the age of the Earth can be measured in billions of years and that the rocks at the surface are the products of a long and complex series of events.

In the late 18th century James Hutton, a Scottish geologist, proposed the classification of rocks into strata or layers. Later, William Smith, who collected fossils in England, noted that some strata contained fossils that were different from those in the strata above and below. This led to the designation of "index fossils" to be used as a basis for dating strata. Figure 4 shows the geologic time scale and compares it to a calendar year. Geologic time has been divided into eras, which are further subdivided into periods and then epochs. The oldest era is the Precambrian, which represents about 80 percent of the total age of the Earth. The Paleozoic (600 to 220 million years ago) is the age of ancient life; the Mesozoic (220 to 70 million years ago) is the age of middle life; and the Cenozoic (70 million years ago to the present) is the age of recent life. The Paleozoic Era has been divided into seven periods of varying lengths. The rocks in Robbers Cave State Park belong to the Pennsylvanian Period, which is toward the end of the Paleozoic Era.

The Pennsylvanian Period is named after the state of Pennsylvania, where rocks of that age contain large amounts of coal. The Pennsylvanian Period began about 320 million years ago and lasted 50 million years. Although periods of geologic time may

seem to be phenomenally long, they are rather brief when compared to the total age of the Earth. When the 4.5-billion-year history of the Earth is compared with a 365-day year, the Pennsylvanian Period began about 26 days ago and lasted only about 4 days (fig. 4).

During the early part of the Pennsylvanian Period the Arkoma Basin (fig. 5), an area of east-central Oklahoma that extends into west-central Arkansas, gradually began to subside. At the same time the region to the south, which is now the Ouachita Mountain Uplift, slowly began to rise.

As these slow movements of the Earth's crustal blocks continued through millions of years, weathering was causing parts of the uplifted Ouachita Mountains to decompose and disintegrate. The sediments produced were carried by streams into the sinking Arkoma Basin, where they were deposited as horizontal layers of strata. After many layers had accumulated in the basin, the sediments were lithified into the solid rocks that can be seen in northern Latimer County and the park area. These sandstones and siltstones have many features similar to present-day delta deposits, such as those in the Mississippi River delta in the New Orleans area. Therefore, using the axiom "the present is the key to the past," geologists who have studied sediments in the park interpret them as being part of an ancient delta deposit.

From time to time during the Pennsylvanian, the Arkoma Basin area was flooded by the shallow marine seas that extended throughout much of the Midcontinent area. The climate during these times was apparently warm and humid, because abundant plant remains suggest that the landscape was covered by luxuriant vegetation. During periods of submergence, marine sediments were deposited, and during periods of emergence, nonmarine sediments were deposited. Thus, interbedded marine and nonmarine sediments extend throughout the basin.

By the end of Pennsylvanian time, more than 21,000 feet of clastic sediments had been deposited in the Arkoma Basin. Geologists believe that the land remained above sea level during the Permian, Triassic, Jurassic, and Cretaceous Periods, because there is no evidence that rocks of these ages were deposited in the Arkoma Basin. Erosion continued through the Tertiary, but alluvium was deposited by streams in the stream valleys during the Quaternary. Because we are still in the Quaternary Period, we can observe that erosion continues to modify the topography, changing features that first formed near the end of the Pennsylvanian Period.

Geologic Time Scale Compared to a Calendar Year

Geologic Era	Geologic Period	Beginning (Million Years Ago)	Comparative Date ¹			
			Day	Hr	Min	
Cenozoic ("Recent life")	Quaternary	1	December	31	22	03
	Tertiary	70	December	26	7	44
Mesozoic ("Middle life")	Cretaceous	135	December	21	1	12
	Jurassic	180	December	17	9	36
	Triassic	220	December	14	3	44
Paleozoic ("Ancient life")	Permian	270	December	10	2	24
	Pennsylvanian	320	December	6	1	04
	Mississippian	350	December	3	14	40
	Devonian	400	November	29	13	20
	Silurian	430	November	27	2	56
	Ordovician	490	November	22	4	37
	Cambrian	600	November	13	16	00
Precambrian		4,500	January	1	0	00

¹Determined by A. J. Myers.

Figure 4. Geologic time scale compared to a calendar year.

Stratigraphy

The branch of geology that deals with descriptions of sedimentary rock, or rock strata, is known as stratigraphy; and examining and describing rocks in the field is the first step for a geologist who is studying an area. The geologist records such things as rock type, color, thickness, orientation, and areal distribution; then from this information he compiles a detailed description of the formations that make up the stratigraphy of the area. From these findings, a geologic map of the area can be produced. Plate 1 (in pocket) is a geologic map of the Robbers Cave State Park area. It shows those parts of T. 6 N. that are present on the Quinton South and Wilburton 7½-minute topographic maps and includes the eastern part of R. 18 E. and most of R. 19 E.

Northern Latimer County was mapped by Dearl T. Russell for his master's thesis in 1958 for the School of Geology and Geophysics at the University of Oklahoma; and Oklahoma Geological Survey Circular 50, *Geology of Northern Latimer County, Oklahoma* (1960), is the published version of that thesis. The stratigraphy of northern Latimer County is summarized by the columnar section in figure 6

that contains the name of each rock unit, lithologic type, and average thickness.

The oldest formation exposed in northern Latimer County is the Atoka Formation. It was named for the town of Atoka in Atoka County, Oklahoma, and is composed of gray to brown silty shale and brown sandstone. The shale is easily eroded and forms valleys, whereas the resistant sandstone forms ridges and caps many hills in the southern part of Latimer County. The Atoka Formation is not exposed in the park or map area.

Overlying the Atoka Formation is the Hartshorne Formation, which is named for the town of Hartshorne in Pittsburg County, Oklahoma. The Hartshorne is gray to brown silty sandstone interbedded with thin, gray silty shales. The formation ranges in thickness from 84 to 316 feet. In the upper part is the upper Hartshorne coal, which is 4 to 6 feet thick and has been mined by surface mining. The Hartshorne Formation is not exposed in the park but is exposed in the extreme southwestern part of the geologic-map area (pl. 1).

The McAlester Formation overlies the Hartshorne Formation and was named for the town of McAlester in Pittsburg County, Oklahoma. It consists of in-




Figure 5. Map showing major geologic provinces of Oklahoma (from Johnson and others, 1972, p. 1). Solid square indicates location of Robbers Cave State Park.

terbedded sandstones and shales and has a total thickness of 2,134 feet. Exposures of the McAlester Formation are in the southwestern and south-central parts of the area mapped (pl. 1). Shales in the road cuts along State Highway 9 are part of the McAlester Formation, along with the coals which have been mined by surface mining. Some of the mines are shown in the south-central part of the map (pl. 1).

The Savanna Formation overlies the McAlester Formation and was named for the town of Savanna, which is also in Pittsburg County, Oklahoma. The Savanna consists of brown, fine-grained sandstone beds and siltstone beds interbedded with gray-green to dark silty shales. There are 14 mappable units in the formation (pl. 1). The sandstones are up to 200 feet thick, and the shales range in thickness from 200 to 550 feet. The total thickness in Latimer County ranges from 1,470 to 1,990 feet.

Most of the rock outcrops in the park are of the Savanna Formation, the most extensive formation in the map area, with many fine exposures in easily accessible road cuts along State Highway 2 and at various localities throughout the park area. Most of the ridges are capped by Savanna sandstones, whereas valleys have formed on the shales. As the shales weather and erode, the thick overlying sandstones form large slump blocks. Some are as much as 30 to 40 feet high. Robbers Cave has formed in sandstone unit IPsv-11 (pl. 1) that also forms the high bluff

upon which cabins have been built. Rough Canyon to the west of Robbers Cave contains large boulders of Savanna sandstones. Shale at the top of the Savanna Formation is exposed along Lake Wayne Wallace and Lake Carlton.

Overlying the Savanna Formation is the Boggy Formation, named from exposures along Boggy Creek in Atoka and Coal Counties, Oklahoma. The Boggy Formation consists of brown to buff sandstones and siltstones interbedded with dark- to light-gray silty shales. There are eight mappable sandstone units with seven interbedded shales in the formation. The Boggy ranges in thickness from 700 feet in western Latimer County to 2,136 feet in eastern Latimer County. The basal sandstone is the Bluejacket Sandstone Member, which forms spectacular bluffs on the west side of Lake Carlton. The Secor coal lies just above the Bluejacket Sandstone. The Boggy Formation is present in the park only in the western part but is present also on the east side of the map area (pl. 1).

Paleontology

Paleontology, the study of life in the geologic past, relies on fossil plants and animals. The fossil




Figure 6. Composite columnar section of rocks in Robbers Cave State Park area. Modified from Russell (1958, pl. 1).

remains of many ancient plants are present in the rocks that crop out in northern Latimer County. Fossil remains of animals are rare, but they can be found with careful searching. Fossil tree stems, stumps, and trunks are common. Sandstone has replaced the original woody material, and imprints of plant leaves are also present in many of the shales—but they are difficult to find. Coal beds in this region nearly always contain the remains of leaves, stems, and spores of many plants.

The most common fossils in the region include *Stigmaria* (fig. 7) and *Calamites* (fig. 8). The two genera of scale trees most common are *Lepidodendron* and *Sigillaria* (fig. 9). In excavating for State Highway 2 many years ago, the thick sandstone layer IPsv-11 of the Savanna Formation (pl. 1), which forms the ridge underlying the park's tourist cabins, yielded the large stump of a scale tree. The road cut where it was found is north of Coon Creek and overlooks Lake Carlton. Specimens of *Stigmaria* are the fossil underground stems of scale trees, and they resemble cacti at first glance. They have been mistaken for the fossil remains of ancient snakes by those not familiar with paleobotany. Numerous *Stigmaria* specimens can be found in creek beds, where they have come to rest after having been eroded out of the strata. *Calamites* specimens, which are numerous, are fossils of a rush-like plant with vertical grooves up the trunk.

Studying fossil plants found in strata helps paleobotanists better understand the Earth's history, because fossils provide information about the conditions and the types of life that existed during the




Figure 7. *Stigmaria*. Fossil underground stem of a scale tree. Pits are where rootlets were attached. These plants are common in the camp area and have been mistaken for fossil cacti or even fossil snakes. From Russell (1958, fig. 11).




Figure 8. *Calamites*. Vertical grooves in the bark characterize this Pennsylvanian rush. Sand casts of this plant are numerous in the park area. From Russell (1958, fig. 10).

time that the enclosing strata were deposited. For example, plant fossils found in the region surrounding the park area are of plants similar to ones that now grow only in warm, humid climates. Therefore, the rock strata in which they are preserved must have been deposited during a period of warm, moist conditions.

Topography and Structural Geology

Topography refers to the shape of the Earth's surface. Robbers Cave State Park is in the Interior Highlands geomorphic province, which includes the Ozark Plateau in northeastern Oklahoma, southwestern Missouri, and northwestern Arkansas, and the Ouachita Mountains in southeastern Oklahoma and west-central Arkansas (fig. 10). The Sans Bois Mountains are in the northwestern part of the Ouachita Mountains, where the region is a sub-mature-to-mature plateau with gently folded strong and weak strata.

The map showing geomorphic provinces of eastern Oklahoma (fig. 11) shows Robbers Cave State Park in the McAlester Marginal Hills Belt. This belt extends from northern Atoka County northeastward




Figure 9. Scale trees. Sandstone casts of the trunks are abundant in the park area. From Russell (1958, fig. 8).

through parts of Coal, Pittsburg, Latimer, and Le Flore Counties. The area is one of resistant Pennsylvanian sandstones that cap broad hills and mountains, rising from 300 to almost 2,000 feet above wide hilly plains.

In the process of mountain building, rocks are often folded into an upward arch (called an anticline) and a downward arch (a syncline). If the deformation is too intense, the strata are broken and are offset as faults. Joints are cracks that form perpendicular to stratification in rocks; commonly two sets of joints occur, in many cases extending at right angles to one another. Weathering and erosion, which result in the removal of rock material, are greater along bedding planes, joints, and faults.

The major structural feature in northern Latimer County is the Sansbois Syncline (pl. 1). The linear extent, or axis, of the fold extends across the middle of the park, with Robbers Cave on the north flank and Old Circle Campground on the south flank, then continues in an east-west direction across the central part of the map area. The topographic highs occur along the axis of the fold. Even though synclines are

downfolds of rock, as erosion continues they commonly form subsequent topographic highs. Figure 12 is a block diagram of a canoe-shaped synclinal mountain west of Robbers Cave State Park, along the axis of the fold. The dashed line is the park boundary.

The North Wilburton Fold (pl. 1) is an anticline shown in the north-central part of the map area. In the northwestern part of the area faults have cut isolated outcrops of the IPsv-11 sandstone member of the Savanna Formation.

The Sans Bois Mountains form the main mountain range between Sans Bois Creek (at an elevation of approximately 550 feet on the north) and Fourche Maline Creek (at an elevation of approximately 600 feet on the south). Blue Mountain in sec. 35, T. 7 N., R. 20 E., northeast of the map area, is the highest point in the Sans Bois Mountains. It is composed of the uppermost sandstone in the Boggy Formation, and the axis of the Sansbois Syncline passes through it. Tucker Knob, elevation 1,532 feet, is a picturesque, isolated, cone-shaped mountain half a mile east of State Highway 2, approximately 7 miles

south of Quinton and 3 miles north of the park (fig. 13). Both Tucker Knob and Panther Mountain, which is 2 miles west of the highway, are composed of the Bluejacket Sandstone, the basal unit of the Boggy Formation.

The topography of the park area consists of high, rugged escarpments or bluffs and enormous blocks of sandstone. The force of gravity has gradually pulled large blocks of sandstone downslope from their original position, and many slump blocks have formed interesting features. In some cases the gradual downslope movement resulted in sandstone layers sliding along smooth bedding planes and giving rise to features such as the steep crevice called Devils Slide.

The high sandstone ridges are separated by valleys that have been cut into the shale layers by streams. Thus the topography of the park area has resulted from selective erosion of rock strata. The height of topographic features is related to the dip (the slope of bedding planes), hardness, and thickness of rock strata. The thicker and more resistant layers form large hills and ridges. Ridges are formed by the folding of the strata forming the Sansbois Syncline, and they are called either hogbacks or cuestas, depending on the slope of the two sides. Hogbacks have slopes that are about the same on both sides; cuestas have one gentle slope and one slope that is much steeper because erosion has cut across the strata. Ridges are well illustrated on the block diagram in figure 12.

Robbers Cave has formed in one of the sandstones of the Savanna Formation (IPsv-11, pl. 1). The sandstone's resistance has caused it to form a cuesta that dips gently to the southwest, into the Sansbois Syncline. On the escarpment side of the cuesta a series of bluffs has formed, attaining a total height of approximately 140 feet (fig. 14). In addition to the folding, the sandstone contains joints and has been faulted. Weathering and erosion have enlarged some of the joints from a few inches to more

than a foot (fig. 15). In some places intermittent streams have sculpted a rounded appearance near the bottom of enlarged joints. On top of the bluff, joints along the bedding plane have the appearance of a checkerboard. Some of the slump blocks have broken away along joint planes (fig. 16). Jointing is responsible for features like the Stone Corral (one of the popular tourist attractions in the park), and some of the narrow passes are also the result of enlarged joint planes.

The sandstone units contain several strata, and smaller cuestas have formed in places on the flanks of the main cuesta (fig. 17). One such cuesta can be seen from near the entrance to Robbers Cave.

The park's tourist cabins have been built along the same cuesta as Robbers Cave but are on the south side of the axis of the Sansbois Syncline, with beds dipping to the northwest. The erosion side of the ridge forms a steep cliff about 200 feet high that descends to Coon Creek, which has formed a valley in the underlying shale (IPsv-10, pl. 1).

The waters of Lake Carlton (fig. 18) cover mainly shale (IPsv-14, pl. 1), but the lake is dammed where Fourche Maline Creek cuts across the sandstone and flows south across other units. The high bluff along the west side of the lake is the erosion side of a cuesta in the Bluejacket Sandstone, which is the basal member of the Boggy Formation.

It is hoped that this guidebook will help visitors to Robbers Cave State Park appreciate the scenic points of interest by encouraging them to look at rocks more closely and to see their makeup. The visitor will gain an even better understanding of geology by thinking about the long period of time and varied environments in which the rocks were formed, and the intense forces that bent the rocks into folds and even caused them to break into faults.

The Earth's surface is not static, and even today dynamic processes are continuing to make changes in Robbers Cave State Park.





Figure 10. Map showing generalized geomorphic regions of Oklahoma and parts of surrounding states. Northeastern Oklahoma is part of the Ozark Plateau, southeastern Oklahoma is part of the Ouachita Mountains, the Panhandle is part of the High Plains, and the rest of the State is part of the Osage Plains of the Central Lowland. Modified from Raisz (1957).



EXPLANATION

Arbuckle Mountain Province



Arbuckle Hills—Low to moderate hills chiefly of Cambrian to Mississippian limestones. Steeply dipping strata are complexly folded and faulted.



Arbuckle Plains—Gently rolling hills and plains developed on granites of Precambrian age and gently dipping limestones chiefly of Ordovician age.

Ouachita Mountain Province



Beavers Bend Hills—Moderate to high hills and ridges formed by tightly folded Ordovician through Mississippian sedimentary rocks.



Ridge and Valley Belt—Long and sinuous mountain ridges of broadly folded Mississippian and Pennsylvanian sandstones towering above subparallel shale valleys.



Hogback Frontal Belt—Thrust blocks of steeply dipping Pennsylvanian sandstones and limestones in hogback ridges rising 500 to 1,700 feet above adjacent valleys.

Other Provinces



Ardmore Basin—Lowland of folded Mississippian and Pennsylvanian shales and sandstones between Arbuckles and Dissected Coastal Plain.



Arkansas Hill and Valley Belt—Broad, gently rolling plains and valleys with scattered hills 100 to 300 feet high capped by Pennsylvanian sandstones.



Boston Mountains—Deeply dissected plateau capped by gently dipping Pennsylvanian sandstones.



Central Redbed Plains—Red Permian shales and sandstones forming gently rolling hills and broad, flat plains.



Claremore Cuesta Plains—Resistant Pennsylvanian sandstones and limestones dipping gently westward, forming cuestas between broad shale plains.



Dissected Coastal Plain—Soft, south-dipping Cretaceous sands, gravels, and clays of Gulf Coastal Plain, slightly dissected by streams.




Eastern Sandstone Cuesta Plains—West-dipping Pennsylvanian sandstones forming cuestas that overlook broad shale plains.



McAlester Marginal Hills Belt—Resistant Pennsylvanian sandstones capping broad hills and mountains that rise 300 to 2,000 feet above wide, hilly plains.



Neosho Lowland—Gently rolling shale lowlands with a few low escarpments and buttes capped by Pennsylvanian sandstones and Mississippian limestones.



Northern Limestone Cuesta Plains—Thin Permian limestones capping west-dipping cuestas that rise above broad shale plains.



Ozark Plateau—Deeply dissected plateau formed in gently dipping Mississippian limestones and cherts.

Figure 11. Map (opposite) showing geomorphic provinces in central and eastern Oklahoma. Solid square indicates location of Robbers Cave State Park. Map adapted from Curtis and Ham (1972).




Figure 12. Schematic aerial view of the canoe-shaped synclinal mountain west of Robbers Cave State Park. The bow of the canoe extends into the park. Dashed line is park boundary. From Russell (1958, fig. 12).




Figure 13a. Tucker Knob is a picturesque, cone-shaped mountain east of State Highway 2 between Quinton and Robbers Cave State Park. Map from U.S. Geological Survey 7½-minute-quadrangle map series: Quinton South, 1969.



Figure 13b. Photograph of Tucker Knob.



Figure 14 (above). Part of the high bluff at Robbers Cave, which is the erosion side of the cuesta formed in sandstone IPsv11 of the Savanna Formation.




Figure 15 (right). One of the many joints that have been enlarged by erosion in sandstone IPsv11 of the Savanna Formation at Robbers Cave. View is at the top of the bluff, with the strata dipping toward the bottom.



Figure 16 (left). Slumped blocks of sandstone result in an almost vertical cliff approximately the size of a three-story building along a joint in the bluff at Robbers Cave.



Figure 17. A small cuesta on the flanks of the main cuesta that has formed in sandstone IPsv11 of the Savanna Formation in the Robbers Cave area. The bedding plane dips toward the left, and on the right is the steep erosional scarp that cuts across the strata.




Figure 18. Lake Carlton and part of the bluff on the erosion side of the cuesta formed by the Bluejacket Sandstone, the basal member of the Boggy Formation.

Botany of Robbers Cave State Park

George J. Goodman¹ and Cheryl A. Lawson²

Introduction

Your visit to Robbers Cave State Park brings you into one of Oklahoma's scenic forests, a forest that lies at the northwestern edge of the Ouachita Mountains and the southern edge of the Sans Bois Mountains. The towering trees, together with the shrubs, vines, and herbaceous plants on the forest floor, blend harmoniously, creating a natural setting for your enjoyment and a sanctuary for the abundant wildlife. The amount of rainfall, the soil type, and other environmental factors are well suited for the development of a forest. If the amount of rainfall were less and the soil different, as they are in central and western Oklahoma, a grassland or prairie would be expected. Even so, in the park there are variations in the makeup of the forest, which is dominated by oaks (*Quercus*) mixed with many hickories (*Carya*) and pines (*Pinus*). On some of the drier slopes and on poor soil, especially where erosion has occurred, the forest is limited to pines. On better soil, mixed with the pines, oaks, and hickories, are several other deciduous species. The most abundant oaks by far on these slopes are post oak (*Q. stellata*) and blackjack (*Q. marilandica*), with the hickories being primarily black (*C. texana*) and mockernut (*C. tomentosa*).

The plants you will find in the park will depend primarily on two factors: the season of your visit and your location in the park. The plants most varied by the season will, of course, be the herbaceous ones. For example, in the spring of the year the forest floor and forest openings may be covered with spring bloomers such as camass (*Camassia scilloides*), crow poison (*Nothoscordum bivalve*), yellow-eyed grass (*Hypoxis hirsuta*), spring beauty (*Claytonia virginica*), buttercups (*Ranunculus*), violet woodsorrel (*Oxalis violacea*), and blue-eyed grass (*Sisyrinchium campestre*), some of which flower before the leafy canopy of the trees develops

and screens out the light. As the season progresses to late summer and fall, very different herbaceous plants will appear. Fewer plants will be seen on the forest floor; however, many will be quite conspicuous with their colors of yellow, blue, and white in the forest openings and along the roadsides. Likely finds are members of the sunflower family (Compositae), such as hairy sunflower (*Helianthus hirsutus*), aster (*Aster ericoides* and *A. sagittifolius*), ironweed (*Vernonia baldwinii*), gayfeather (*Liatris squarrosa*), broomweed (*Gutierrezia dracunculoides*), plains coreopsis (*Coreopsis tinctoria*), goldenrod (*Solidago*), and fragrant cudweed (*Gnaphalium obtusifolium*).

It is true, too, that certain trees and shrubs will be more conspicuous than others at certain times of the year. For example, flowering dogwood (*Cornus florida*) with its beautiful flower-like bracts and redbud (*Cercis canadensis*), our state tree, with its abundant pinkish-red flowers can be viewed easily in the early spring through what remains a predominantly dormant forest. In many parts of the park, French mulberry (*Callicarpa americana*) adds a splash of fall color with its wand-like stems covered in clusters of purple berries. This is one of Oklahoma's truly lovely shrubs.

Slopes, particularly north-facing ones, and protected ravines are more moist areas and better support the growth of plants requiring more moisture. The trees you may find in this area include chinquapin oak (*Q. muhlenbergii*), redbud (*Cercis*), flowering dogwood (*C. florida*), spotted oak (*Q. shumardii*), red maple (*Acer rubrum*), and winged elm (*Ulmus alata*). Occasional are hackberry (*Celtis*), willow oak (*Q. phellos*), black cherry (*Prunus serotina*), and chittamwood (*Bumelia lanuginosa*).

Likely shrubs to be found in these more moist areas are witch hazel (*Hamamelis vernalis*), black-

¹Regents Professor Emeritus, Department of Botany and Microbiology, University of Oklahoma, Norman.

²Adjunct assistant professor, Department of Botany and Microbiology, University of Oklahoma, Norman.

berry (*Rubus*), prairie rose (*Rosa setigera*), dense St. Johns wort (*Hypericum densiflorum*), falseindigo (*Amorpha fruticosa*), and buttonbush (*Cephalanthus occidentalis*).

Along Fourche Maline Creek, black willow (*Salix nigra*), river birch (*Betula nigra*), American elm (*U. americana*), sycamore (*Platanus occidentalis*), and various oaks are characteristic. Mixed with the trees and shrubs in the summer are such herbs as brown-eyed Susan (*Rudbeckia triloba*) and mistflower (*Eupatorium coelestinum*).

At certain times of the year, Lake Carlton itself may support an abundance of aquatic plants, such as pondweed (*Potamogeton*), bulrush (*Scirpus*), Brazilian parrotfeather (*Myriophyllum brasiliense*), water-willow (*Justicia americana*), and water hyssop (*Bacopa rotundifolium*). Frequent on the lake bor-

ders are cat-tail (*Typha domingensis*), water hemlock (*Cicuta maculata*), small-spike false-nettle (*Boehmeria cylindrica*), and rush (*Juncus*).

Roadsides and prairie openings support the growth of many grasses along with tree huckleberry (*Vaccinium arboreum*); smooth, winged, and fragrant sumac (*Rhus glabra*, *R. copallina*, and *R. aromatica*); aster (*Aster*); goldenrod (*Solidago*); and bush-clover (*Lespedeza*). In the fall such grasses as silver bluestem (*Andropogon saccharoides*), big bluestem (*A. gerardii*), little bluestem (*A. scoparius*), long-spike tridens (*Tridens strictus*), Indian grass (*Sorghastrum nutans*), and Johnson grass (*Sorghum halapense*) can be expected in one or both of these areas. At the forest edge, coralberry (*Symphoricarpos orbiculatus*) occurs often.

Key to the Woody Plants

The following key to the woody plants of the park and adjacent areas is based almost entirely on leaf and twig characters. Hence, reasonably mature leaves should be used.

Meanings of terms, such as "opposite," "alternate," "simple," and "compound," can be ascertained from the illustrations in figures 19 and 20. A ruler is provided at the edge of the inside back cover for measurements needed in the key.

In using the key, you will find that there are always two choices. These are indicated both by indentation and by paired symbols (A and AA, B and

BB, etc.). Generally, it is better to read both choices, then choose the better one. As an example, suppose we had a Spotted Oak (fig. 21) which we wish to identify. The leaves are not needle- or scale-like (as described at A), so we choose AA. The leaves are arranged alternately on the stem (one at a node), so we would choose C not CC, where the leaves are to be opposite (two at a node). The leaves are simple, so choose D rather than DD. But the leaves are not entire; instead they are lobed, so we pass E and go to EE, where it states that the leaves are toothed or lobed. The plant is not a vine but a tree, so choose SS. The leaves are lobed halfway or more to the middle, so we choose V. Soon you are at Spotted Oak. Good luck! Have fun!

- A. Trees evergreen and with leaves either scale-like and small or needle-like.
 - B. Leaves scale-like, most of them hardly an eighth of an inch long; fruits berry-like Eastern red cedar (*Juniperus virginiana*)
 - BB. Leaves needle-like, 2 inches or more long. Short leaf pine (*Pinus echinata*; fig. 22)
- AA. Trees, shrubs or vines, but not with leaves as above.
 - C. Leaves alternate.
 - D. Leaves simple.
 - E. Leaves entire.
 - F. Leaves about as long as wide, or at least less than twice as long as wide.
 - G. Vines.
 - H. Stems prickly. Greenbrier (*Smilax*)
 - HH. Stems not prickly. Red berry moonseed (*Cocculus carolinus*)
 - GG. Trees or shrubs.
 - I. Veins 5 or more from the base of the heart-shaped leaf. Redbud (*Cercis canadensis*; fig. 24)
 - II. Veins 3 from the lop-sided leaf base. Georgia hackberry (*Celtis tenuifolia*; fig. 25)
 - FF. Leaves 2 or more times longer than wide.
 - J. Woody vines. Bama supplejack (*Berchemia scandens*)
 - JJ. Trees or shrubs.
 - K. Trees.
 - L. Leaves tipped with a bristle an eighth of an inch or more long and the leaf stalk less than 0.5 inch long Willow oak (*Quercus phellos*)
 - LL. Leaves without bristle tips, or if bristle tipped, the leaf stalk an inch or more long.
 - M. Leaves with 3 major veins (best seen from beneath) arising from the base of the lop-sided leaf. Sugar hackberry (*Celtis laevigata*)
 - MM. Leaves with several veins arising along the length of the midrib.
 - N. Larger leaf blades over 8 inches long ... Pawpaw (*Asimina triloba*)
 - NN. Leaf blades not over 4 inches long.
 - O. End of leaves rounded ... Chittamwood (*Bumelia lanuginosa*)
 - OO. End of leaves with a distinct point or tip.
 - P. Leaves glossy, the twigs often with thorns about 0.5 inch long Osage orange (*Maclura pomifera*)
 - PP. Leaves not glossy and twigs without thorns. Persimmon (*Diospyros virginiana*)
 - Black tupelo (*Nyssa sylvatica*)
(These two are most easily separated by the fruits, those of persimmon being a nearly spherical berry often over an inch in diameter, and those of black tupelo being elongate fleshy fruits 0.5 inch long).
 - KK. Shrubs.
 - Q. Leaves glossy, thick and leathery; tall shrubs. Tree huckleberry (*Vaccinium arboreum*; fig 23)
 - QQ. Leaves not glossy or thick; plants usually not over 4 feet tall.
 - R. Leaves very often rounded at the base Deerberry (*Vaccinium stamineum*)
 - RR. Leaves tapering to the base; plants commonly not over 2 feet tall. Blueridge blueberry (*Vaccinium vacillans*)
 - EE. Leaves toothed or lobed (sometimes the lobes or teeth only near the end of the leaf).
 - S. Vines.

- T. Stems prickly Greenbrier (*Smilax*)
- TT. Stems not prickly.
 - U. Flower and fruit clusters longer than wide Grape (*Vitis*)
 - UU. Flower and fruit clusters wider than long Heart leaf ampelopsis (*Ampelopsis cordata*)
- SS. Trees or shrubs.
 - V. Leaves lobed half way or more to the middle.
 - W. Lobes ending in a slender bristle.
 - X. Under surface of leaves hairless Spotted oak (*Quercus shumardii*; fig. 21)
 - XX. Under surface of leaves short-hairy Black oak (*Quercus velutina*)
 - WW. Lobes not ending in a bristle.
 - Y. Leaves with teeth as well as lobes; occasional leaves of Red Mulberry (*Morus rubra*; fig. 23)
 - YY. Leaves lobed but without teeth Post oak (*Quercus stellata*; fig. 22)
 - VV. Leaves toothed or wavy margined but not deeply lobed.
 - Z. Leaves large (commonly 5 inches or more wide), about as broad as long, angularly lobed American sycamore (*Platanus occidentalis*; fig. 23)
 - ZZ. Leaves more nearly triangular, oval, or narrow or obviously broadest above the middle.
 - a. Leaves broadest above the middle.
 - b. Leaves mostly 3 to 4 inches long.
 - c. Widest portion of the leaves about 1.5 inches Water oak (*Quercus nigra*; fig. 21)
 - cc. Widest portion of the leaves 2.5 inches or more Blackjack oak (*Quercus marilandica*; fig. 21)
 - bb. Leaves mostly 2.5 inches or less long.
 - d. Leaves broadest well above the middle; twigs sometimes thorny; petals 5, in loose clusters at the end of branchlets; fruits with remaining flower parts at the top as in an apple Little hip hawthorne (*Crataegus spathulata*)
 - dd. Leaves broadest only slightly above the middle; twigs never thorny; petals 4, flowers mostly single or few in a cluster on short stalks along the stem; fruits with remaining flower parts, if any, at the stem end more nearly like a cherry Possumhaw (*Ilex decidua*)
 - aa. Leaves broadest near the middle or below.
 - e. Leaves broadest near their lop-sided base; 3 major veins arising from the leaf base; leaves 2 to 4 or more inches long, alternating on but two sides of the twigs Hackberry (*Celtis occidentalis*)
 - ee. Leaves differing in one or more ways from the above.
 - f. Leaf stalks commonly 2 inches or more long, as long or longer than the triangular leaf Eastern cottonwood (*Populus deltoides*; fig. 25)
 - ff. Leaf stalks less than 2 inches long, never so long as the leaf.
 - g. Base of leaf on each side of midrib straight and parallel with the straight lateral veins River birch (*Betula nigra*; fig. 22)
 - gg. Base of leaf rounded, tapering, or heartshaped, hence if lateral veins straight, then not parallel to the leaf base.
 - h. Leaves more than 3 inches long.
 - i. Leaves 0.5 inch or less wide Black willow (*Salix nigra*; fig. 22)
 - ii. Leaves an inch or more wide.
 - j. Teeth over 20 on each edge of the leaf.
 - k. Veins on each side of the midrib fewer than 10 Red mulberry (*Morus rubra*; fig. 23)
 - kk. Veins on each side of the midrib more than 10 American elm (*Ulmus americana*; fig. 24)
 - jj. Teeth fewer than 15 on each edge of the leaf; leaves sometimes appearing as wavy margined.
 - l. Veins about 10 on each side of the midrib Chinquapin oak (*Quercus muhlenbergii*; fig. 21)
 - ll. Veins about 6 on each side of the midrib; leaves often but wavy margined Ozark witch-hazel (*Hamamelis vernalis*; fig. 24)
 - hh. Leaves 3 inches or less long.
 - m. Leaves about as broad as long, broadly triangular to nearly round in outline, sharply toothed; twigs usually thorny ... Downy hawthorne (*Crataegus mollis*)


- mm. Leaves at least twice as long as wide; twigs not thorny.
- n. Shrubs not over 4 feet tall.
 - o. Leaves with 3 evident veins from the base.
 - p. Leaves at least 2 inches long Jerseytea (*Ceanothus americanus*)
 - pp. Leaves less than 2 inches long Inland ceanothus (*Ceanothus herbaceus*)
 - oo. Leaves with veins arising along the midrib; teeth on edges of the leaf very small Blueridge blueberry (*Vaccinium vacillans*)
- nn. Trees, or if shrubs, over 6 feet tall.
 - q. Leaf stalks less than 0.25 inch long; trees.
 - r. Leaves about 2 inches long; twigs usually winged Winged elm (*Ulmus alata*; fig. 25)
 - rr. Leaves nearer 4 inches long; twigs not winged American elm (*Ulmus americana*; fig. 24)
 - qq. Leaf stalks at least 0.25 inch long, usually longer.
 - s. Leaves, at least mostly, less than an inch wide.
 - t. Leaves slender, 3 times or more as long as wide; often tree-sized shrubs with several trunks from the base Black willow (*Salix nigra*; fig. 22)
 - tt. Leaves usually no more than twice as long as wide, widest at the middle or above Possumhaw (*Ilex decidua*)
 - ss. Leaves, at least mostly, an inch or more wide.
 - u. Leaf bases rounded or tapering, leaf stalks rarely as much as 0.75 inch long, mostly with one or two glands near attachment to the leaf blade; remnants of the flower at the stem end of the fruit; flowers appearing with the leaves. Black cherry (*Prunus serotina*)
 - uu. Leaf bases frequently heart-shaped, leaf stalks often 0.75 to 1 inch long and lacking glands; remnants of the flower at the top of the fruit as in an apple; flowers with their long, narrow (0.5 to 0.75 inch long) petals appearing before the leaves Serviceberry (*Amelanchier arborea*)

DD. Leaves compound.

- v. Leaflets 3 to 5, all, except sometimes the middle one, arising from one point.
 - w. Leaflets 5; a climbing, non-spiny vine Virginia creeper (*Parthenocissus quinquefolia*)
 - ww. Leaflets 3, or if sometimes 5, the stems prickly; shrubs or vines.
 - x. Stems lacking spines or prickles.
 - y. Leaflets 2 to 3 inches long, margins toothed Fragrant sumac (*Rhus aromatica*; fig. 28)
 - yy. Leaflets to 4 inches long, margins nearly smooth to irregularly toothed. Poison ivy (*Rhus radicans*; fig. 28)

- xx. Stems spiny or prickly; shrubs.
 - z. Leaf stalks prickly. Blackberry & Dewberry (*Rubus*; fig. 26)
 - zz. Leaf stalks lacking prickles; leaflets mostly 3 Prairie rose (*Rosa setigera*; fig. 27)
- vv. Leaflets often more than 5, and not arising from one point.
 - 1. Shrubs, usually not over 8 feet tall.
 - 2. Midrib of the leaf winged. Winged sumac (*Rhus copallina*; fig. 28)
 - 2'. Midrib of leaf wingless.
 - 3. Leaflets toothed.
 - 4. Stems prickly Leafy rose (*Rosa foliolosa*)
 - 4'. Stems not prickly Smooth sumac (*Rhus glabra*; fig. 26)
 - 3'. Leaflets entire or nearly so Falseindigo (*Amorpha fruticosa*)
 - 1'. Trees.
 - 5. Leaflets 2 inches long or less, entire or minutely wavy margined.
 - 6. Most of the leaflets more than 0.5 inch wide Black locust (*Robinia pseudoacacia*)
 - 6'. Most of the leaflets less than 0.5 inch wide. Honey locust (*Gleditsia triacanthos*)
 - 5'. Leaflets over 2 inches long, toothed.
 - 7. Leaflets 9 or fewer Hickory (*Carya*; fig. 27)
 - 7'. Leaflets 11 or more Black walnut (*Juglans nigra*)
- CC. Leaves opposite, (sometimes 3 to 4 at a node).
 - 8. Leaves simple.
 - 9. Leaves entire.
 - 10. Vines.
 - 11. Leaves whitish beneath Yellow honeysuckle (*Lonicera flava*)
 - 11'. Leaves green beneath American starjasmine (*Trachelospermum difforme*)
 - 10'. Upright trees or shrubs.
 - 12. Leaves often 3 to 4 at a node; flowers and fruits in spherical clusters Buttonbush (*Cephalanthus occidentalis*; fig. 29)
 - 12'. Leaves 2 per node.
 - 13. Shrubs usually less than 4 feet tall, or, if sometimes a little taller, the leaves no more than 2 inches long.
 - 14. Leaves twice or less as long as wide Coralberry (*Symphoricarpos orbiculatus*; fig. 29)
 - 14'. Leaves more than twice as long as wide Dense St. Johns wort (*Hypericum densiflorum*)
 - 13'. Shrubs or small trees more than 4 feet tall.
 - 15. Flowers and fruits in dense clusters; petal-like bracts which appear in the spring large and white or sometimes pink. Flowering dogwood (*Cornus florida*; fig. 29)
 - 15'. Flowers and fruits in loose clusters often 2 or 3 inches wide and often as long.
 - 16. Leaves 1.5 inches or more wide, lower surface only slightly paler green than the upper Rough leaf dogwood (*Cornus drummondii*)
 - 16'. Leaves frequently less than 1.5 inches wide, lower surface much paler green than the upper, appearing whitish Silky dogwood (*Cornus obliqua*)
 - 9'. Leaves toothed or lobed.
 - 17. Leaves lobed and up to 2 inches long; plant a low shrub; young shoots of Coralberry (*Symphoricarpos orbiculatus*; fig. 29)
 - 17'. Leaves toothed and more than 2 inches long.
 - 18. Leaves less than twice as long as wide Red maple (*Acer rubrum*; fig. 29)
 - 18'. Leaves more than twice as long as wide.
 - 19. Leaves usually 4 to 6 inches long; flowers and fruits in clusters at the base of the leaves French mulberry (*Callicarpa americana*; fig. 29)
 - 19'. Leaves less than 4 inches long; flowers and fruits not arranged as in the preceding.
 - 20. Leaves glossy, the leaf stalks very usually covered with rusty red hairs; flowers and fruits borne at the ends of the twigs Southern blackhaw (*Viburnum rufidulum*)
 - 20'. Leaves not glossy; flowers and fruits borne at the leaf bases on stalks an inch or more long Eastern wahoo (*Euonymus atropurpureus*)
 - 8'. Leaves compound.
 - 21. Vines.
 - 22. Leaflets toothed; stems woody; flowers red Common trumpetvine (*Campsis radicans*)
 - 22'. Leaflets entire or sometimes lobed; stems hardly woody; flowers bluish Leatherflower (*Clematis pitcheri*)
 - 21'. Upright trees or shrubs.
 - 23. Leaflets 3, only occasionally 5 Boxelder (*Acer negundo*)
 - 23'. Leaflets 5 to 11.
 - 24. Leaflets entire or with minute teeth; trees Ash (*Fraxinus*)
 - 24'. Leaflets sharply toothed; shrubs American elder (*Sambucus canadensis*)

Leaf Organization



Pinnately compound






Figure 19. Illustrations showing leaf organization.

Leaf Parts



Leaf Position



Leaf Margins






Figure 20. Illustrations showing leaf parts, position, and margins.




Blackjack oak
(*Quercus marilandica*)



Spotted oak
(*Quercus shumardii*)



Chinquapin oak
(*Quercus muhlenbergii*)



Water oak
(*Quercus nigra*)

Figure 21. Illustrations showing simple, alternate leaved species.







Figure 22. Illustrations showing simple, alternate leaved species.




Tree huckleberry
(*Vaccinium arboreum*)



Red mulberry
(*Morus rubra*)



Red mulberry
(*Morus rubra*)



American sycamore
(*Platanus occidentalis*)

Figure 23. Illustrations showing simple, alternate leaved species.




Figure 24. Illustrations showing simple, alternate leaved species.




Figure 25. Illustrations showing simple, alternate leaved species.




Figure 26. Illustrations showing compound, alternate leaved species.




Figure 27. Illustrations showing compound, alternate leaved species.





Figure 28. Illustrations showing compound, alternate leaved species.



List of the Plants

The following is a list of plants from Robbers Cave State Park and the area surrounding it. Nearly all the plants on this list are to be expected within

the park; however, only those marked with an asterisk have been recorded within its boundaries.

The names occurring on the left side of each column are the scientific names. The names to the right are the common names.

POLYPODIACEAE

- **Asplenium pinnatifidum*
- **Asplenium platyneuron*
- Cheilanthes lanosa*
- **Cheilanthes tomentosa*
- Pellaea atropurpurea*
- **Polypodium polypodioides*
- **Pteridium aquilinum*
- Woodsia obtusa*

PINACEAE

- **Juniperus virginiana*
- **Pinus echinata*

TYPHACEAE

- **Typha domingensis*

POTAMOGETONACEAE

- **Potamogeton*

ALISMACEAE

- Alisma subcordatum*
- Sagittaria ambigua*

GRAMINEAE

- **Agrostis elliottiana*
- **Andropogon gerardii*
- **Andropogon saccharoides*
- **Andropogon scoparius*
- **Andropogon virginicus*
- **Aristida oligantha*
- Arundinaria gigantea*
- **Bromus japonicus*
- **Bromus mollis*
- **Cynodon dactylon*
- **Digitaria sanguinalis*
- **Elymus canadensis*
- Eragrostis capillaris*
- **Eragrostis spectabilis*
- Manisuris cylindrica*
- **Panicum anceps*
- **Panicum dichotomum*
- Panicum hians*
- **Panicum lanuginosum* var. *fasciculatum*
- **Panicum malacophyllum*
- **Panicum oligosanthos* var. *scribnerianum*
- Panicum sphaerocarpon*
- **Paspalum dilatatum*
- **Paspalum floridanum* var. *floridanum*
- **Setaria glauca*
- **Sorghastrum nutans*
- **Sorghum halapense*
- **Sphenopholis obtusata*
- **Sporobolus asper*

FERN FAMILY

- Pinnatifid spleenwort
- Brown stem spleenwort
- Hairy lipfern
- Woolly lipfern
- Purple cliffbrake
- Resurrection fern
- Western bracken
- Blunt lobe cliff fern

PINE FAMILY

- Eastern redcedar
- Short leaf pine

CAT-TAIL FAMILY

- Narrow leaf cat-tail

PONDWEED FAMILY

- Pondweed

WATER-PLANTAIN FAMILY

- Small flower water-plantain
- Arrowhead

GRASS FAMILY

- Annual ticklegrass
- Big bluestem
- Silver bluestem
- Little bluestem
- Yellow sedge bluestem
- Prairie threeawn
- Giant cane
- Japanese brome
- Soft brome
- Bermuda grass
- Hairy crabgrass
- Canada wildrye
- Lacegrass
- Purple lovegrass
- Carolina joint-tail
- Beaked panicum
- Forked panicum
- Gaping panicum

- Woolly panicum
- Soft leaf panicum

- Scribner panicum
- Round seed panicum
- Dallisgrass

- Florida paspalum
- Foxtail
- Indian grass
- Johnson grass
- Prairie wedgescale
- Tall dropseed

GRAMINEAE (continued)

- **Tridens flavus*
- **Tridens strictus*
- **Uniola latifolia*

CYPERACEAE

- Carex annectens*
- **Carex austrina*
- **Carex blanda*
- **Carex bushii*
- Carex crus-corvi*
- **Carex emoryi*
- Carex frankii*
- **Carex louisianica*
- Carex lupulina*
- Carex lurida*
- Carex squarrosa*
- Carex straminea*
- Cyperus ovularis*
- Cyperus strigosus*
- Rhynchospora globularis* var.
harveyi
- **Scirpus*

COMMELINACEAE

- Commelina virginica*
- Tradescantia ohiensis*

JUNCACEAE

- **Juncus acuminatus*
- Juncus brachycarpus*
- Juncus diffusissimus*
- **Juncus effusus*
- **Juncus interior*
- **Juncus marginatus*
- Juncus validus*
- **Luzula campestris* var.
bulbosa

LILIACEAE

- **Camassia scilloides*
- **Nothoscordum bivalve*
- **Smilax bona-nox*
- Smilax glauca*
- Smilax tamnoides*
- **Zygadenus nuttallii*

AMARYLLIDACEAE

- Agave lata*
- **Cooperia drummondii*
- **Hypoxis hirsuta*

IRIDACEAE

- **Sisyrinchium campestre*

ORCHIDACEAE

- **Spiranthes gracilis*
- Spiranthes vernalis*

SALICACEAE

- **Salix nigra*
- **Populus deltoides*

JUGLANDACEAE

- Carya cordiformis*
- **Carya texana*
- **Carya tomentosa*
- Juglans nigra*

GRASS FAMILY (continued)

- Purpletop
- Long spike tridens
- Broad leaf spanglegrass

SEDGE FAMILY

- Yellow fruit sedge
- Southern sedge
- Woodland sedge
- Bush sedge
- Crowfoot sedge
- Emory sedge
- Franks sedge
- Louisiana sedge
- Hop sedge
- Lurid sedge
- Squarrose sedge
- Straw sedge
- Globe flatsedge
- False nutgrass

- Globe beakrush
- Bulrush

SPIDERWORT FAMILY

- Virginia dayflower
- Ohio spiderwort

RUSH FAMILY

- Knot leaf rush
- White root rush
- Slimpod rush
- Soft rush
- Inland rush
- Grass leaf rush
- Round head rush

- Bulb woodrush

LILY FAMILY

- Eastern camass
- Crow poison
- Saw greenbrier
- Cat greenbrier
- Bristly greenbrier
- Poison camass

AMARYLLIS FAMILY

- Agave
- Evening star rainlily
- Yellow-eyed grass

IRIS FAMILY

- Blue-eyed grass

ORCHID FAMILY

- Slender ladiestresses
- Upland ladiestresses

WILLOW FAMILY

- Black willow
- Eastern cottonwood

WALNUT FAMILY

- Bitternut hickory
- Black hickory
- Mockernut hickory
- Black walnut

BETULACEAE

**Alnus serrulata*

**Betula nigra*

FAGACEAE

Quercus alba

**Quercus borealis*

Quercus falcata

**Quercus lyrata*

Quercus macrocarpa

**Quercus margaretta*

**Quercus marilandica*

**Quercus muhlenbergii*

**Quercus nigra*

**Quercus phellos*

**Quercus shumardii*

**Quercus stellata*

**Quercus velutina*

ULMACEAE

Celtis laevigata

Celtis occidentalis

**Celtis tenuifolia*

**Ulmus alata*

**Ulmus americana*

**Ulmus crassifolia*

Ulmus rubra

MORACEAE

**Maclura pomifera*

**Morus alba*

**Morus rubra*

URTICACEAE

**Boehmeria cylindrica*

LORANTHACEAE

**Phoradendron serotinum*

POLYGONACEAE

Polygonum bicornis

Polygonum convolvulus

Polygonum hydropiper

Polygonum opelousanum

**Polygonum pennsylvanicum*

Polygonum persicaria

Polygonum tenue

Polygonum virginianum

Rumex hastatulus

Rumex verticillatus

NYCTAGINACEAE

**Mirabilis albida*

PHYTOLACCACEAE

**Phytolacca americana*

AIZOACEAE

Mollugo verticillata

PORTULACACEAE

**Claytonia virginica*

CARYOPHYLLACEAE

**Arenaria drummondii*

**Arenaria patula* var.

robusta

**Cerastium brachypodium*

**Cerastium vulgatum*

Holosteum umbellatum

BIRCH FAMILY

Alder

River birch

BEECH FAMILY

White oak

Northern red oak

Southern red oak

Overcup oak

Bur oak

Dwarf post oak

Blackjack oak

Chinquapin oak

Water oak

Willow oak

Spotted oak

Post oak

Black oak

ELM FAMILY

Sugar hackberry

Hackberry

Georgia hackberry

Winged elm

American elm

Cedar elm

Red elm

MULBERRY FAMILY

Osage orange

White mulberry

Red mulberry

NETTLE FAMILY

Small spike false nettle

MISTLETOE FAMILY

Christmas mistletoe

BUCKWHEAT FAMILY

Smartweed

Dull seed cornbind

Smartweed

Smartweed

Pennsylvania smartweed

Spotted smartweed

Pleat leaf knotweed

Virginia knotweed

Heartwing sorrell

Swamp dock

FOUR-O'CLOCK FAMILY

White four-o'clock

POKEWEED FAMILY

Pokeweed

CARPETWEED FAMILY

Green carpetweed

PURSLANE FAMILY

Spring beauty

PINK FAMILY

Drummond sandwort

Pitcher sandwort

Short stalk chickweed

Big cerastium

Jagged chickweed

**Silene antirrhina*
Silene stellata
RANUNCULACEAE
Clematis pitcheri
Delphinium carolinianum
Delphinium virescens
**Ranunculus fascicularis*
Ranunculus laxicaulis
**Ranunculus macranthus*
BERBERIDACEAE
**Podophyllum peltatum*
MENISPERMACEAE
**Cocculus carolinus*
**Menispermum canadense*
ANONACEAE
Asimina triloba
LAURACEAE
Sassafras albidum var.
molle
FUMARIACEAE
**Corydalis crystallina*
CRUCIFERAE
**Arabis canadensis*
**Brassica kaber*
**Lepidium virginicum*
**Sisymbrium officinale*
SAXIFRAGACEAE
Penthorum sedoides
HAMAMELIDACEAE
**Hamamelis vernalis*
PLATANACEAE
**Platanus occidentalis*
ROSACEAE
**Amelanchier arborea*
**Crataegus mollis*
Crataegus spathulata
Fragaria virginiana var.
illinoensis
Geum canadense
Gillenia stipulata
**Potentilla simplex*
**Prunus serotina*
Rosa carolina
Rosa foliolosa
**Rosa setigera*
**Rubus ostryifolius*
**Rubus trivialis*
LEGUMINOSAE
**Amorpha fruticosa*
**Apios americana*
Astragalus distortus
**Astragalus mexicanus*
Baptisia leucantha
**Baptisia leucophaea*
**Baptisia sphaerocarpa*
**Cassia chamaecrista*
**Cercis canadensis*
**Clitoria mariana*
Crotalaria sagittalis

Sleepy catchfly
 Starry campion
BUTTERCUP FAMILY
 Leatherflower
 Blue larkspur
 Plains larkspur
 Tufted buttercup
 Water plantain spearwort
 Showy buttercup
BARBERRY FAMILY
 May-apple
MOONSEED FAMILY
 Red berry moonseed
 Canada moonseed
CUSTARD-APPLE FAMILY
 Pawpaw
LAUREL FAMILY

 Sassafras
FUMATORY FAMILY
 Mealy corydalis
MUSTARD FAMILY
 Sicklepod
 Charlock
 Poor man's pepper
 Hedgemustard
SAXIFRAGE FAMILY
 Virginia penthorum
WITCH-HAZEL FAMILY
 Ozark witch-hazel
PLANE-TREE FAMILY
 American sycamore
ROSE FAMILY
 Serviceberry
 Downy hawthorne
 Little hip hawthorne

 Wild strawberry
 White avens
 Indian physic
 Old field cinquefoil
 Black cherry
 Carolina rose
 Leafy rose
 Prairie rose
 High brush blackberry
 Southern dewberry
BEAN FAMILY
 Falseindigo
 American potatobean
 Bent pod milkvetch
 Ground plum
 White falseindigo
 Plains wildindigo
 Yellow wildindigo
 Prairie-senna
 Redbud
 Atlantic pigeon wings
 Arrow crotalaria

Desmodium canescens
 **Desmodium ciliare*
Desmodium dillenii
Desmodium rigidum
Desmodium sessilifolium
 **Galactia volubilis*
 **Gleditsia triacanthos*
Lathyrus pusillus
 **Lespedeza cuneata*
Lespedeza repens
 **Lespedeza stipulacea*
 **Lespedeza stuevei*
Lespedeza virginica
 **Melilotus alba*
Neptunia lutea
 **Petalostemon candidum*
Psoralea floribunda
 **Psoralea psoralioides* var.
 eglandulosa
Rhynchosia latifolia
 **Robinia pseudoacacia*
 **Schrankia nuttallii*
 **Strophostyles leiosperma*
 **Strophostyles umbellata*
 **Stylosanthes biflora*
 **Tephrosia virginiana*
Trifolium campestre
 **Trifolium carolinianum*
 **Trifolium dubium*
 **Trifolium reflexum*

OXALIDACEAE

**Oxalis corniculata*
Oxalis stricta
 **Oxalis violacea*

POLYGALACEAE

**Polygala incarnata*
Polygala sanguinea
 **Polygala verticillata* var.
 dolichoptera

EUPHORBIACEAE

**Acalypha gracilens* var.
 monococca
Acalypha rhomboidea
Andrachne phyllanthoides
 **Croton capitatus*
 **Croton monanthogynus*
Crotonopsis elliptica
Euphorbia corollata
 **Euphorbia dentata*
Euphorbia spathulata
 **Euphorbia supina*
 **Tragia ramosa*

CALLITRICHACEAE

**Callitriche heterophylla*

ANACARDIACEAE

**Rhus aromatica*
 **Rhus copallina*
 **Rhus glabra*
 **Rhus radicans*
 **Rhus toxicodendron*

Hoary tickclover
 Little leaf tickclover
 Panicked tickclover
 Stiff tickclover
 Sessile leaf tickclover
 Downy milkpea
 Honeylocust
 Low peavine
 Chinese bushclover
 Creeping bushclover
 Korean bushclover
 Stueves bushclover
 Slender bushclover
 White sweetclover
 Yellow neptunia
 White prairieclover
 Many flower psoralea

Sampson snakeroot
 Broad leaf snoutbean
 Black locust
 Nuttall sensitive-brier
 Slick seed wildbean
 Perennial wildbean
 Pencil flower
 Devil's shoestring
 Low hop clover
 Carolina clover
 Shamrock
 Buffalo clover

WOODSORREL FAMILY

Creeping oxalis
 Yellow oxalis
 Violet woodsorrel

MILKWORT FAMILY

Pink milkwort
 Blood milkwort

Whorled milkwort

SPURGE FAMILY

Slender copperleaf
 Rhomboid copperleaf
 Missouri maidenbush
 Woolly croton
 Single fruit croton
 Narrow leaf rushfoil
 Flowering spurge
 Toothed spurge
 Warty euphorbia
 Spotted euphorbia
 Noseburner

WATER-STARWORT FAMILY

Water-starwort

CASHEW FAMILY

Fragrant sumac
 Winged sumac
 Smooth sumac
 Poison ivy
 Poison oak

AQUIFOLIACEAE

**Ilex decidua*

CELASTRACEAE

Euonymus atropurpureus

ACERACEAE

Acer negundo

**Acer rubrum*

Acer saccharinum

BALSAMINACEAE

Impatiens capensis

RHAMNACEAE

Berchemia scandens

Ceanothus americanus var.
pitcheri

**Ceanothus herbaceus* var.
pubescens

**Rhamnus caroliniana*

VITACEAE

Ampelopsis cordata

**Cissus incisa*

**Parthenocissus quinquefolia*

**Vitis* sp.

MALVACEAE

Callirhoe alcaeoides

Sida spinosa

HYPERICACEAE

**Ascyrum hypericoides* var.
multicaule

**Hypericum densiflorum*

**Hypericum drummondii*

Hypericum mutilum

**Hypericum punctatum*

CISTACEAE

**Lechea tenuifolia*

VIOLACEAE

Viola missouriensis

**Viola pedata*

**Viola pennsylvanica*

**Viola rafinesquii*

PASSIFLORACEAE

**Passiflora incarnata*

Passiflora lutea var.
glabriflora

NYSSACEAE

**Nyssa sylvatica*

MELASTOMACEAE

**Rhexia mariana* var.
interior

ONAGRACEAE

**Gaura longiflora*

**Ludwigia alternifolia* var.
pubescens

Oenothera biennis

**Oenothera fruticosa* var.
Goodmanii

**Oenothera laciniata*

Oenothera speciosa

HALORAGIDACEAE

**Myriophyllum brasiliense*

HOLLY FAMILY

Possumhaw

STAFF-TREE FAMILY

Eastern wahoo

MAPLE FAMILY

Boxelder

Red maple

Silver maple

TOUCH-ME-NOT FAMILY

Spotted snapweed

BUCKTHORN FAMILY

Bama supplejack

Jerseytea

Inland ceanothus

Carolina buckthorn

GRAPE FAMILY

Heart leaf ampelopsis

Ivy treebine

Virginia creeper

Grape

MALLOW FAMILY

Plains poppymallow

Prickly mallow

ST. JOHNS WORT FAMILY

St. Andrews cross

Dense St. Johns wort

Drummond St. Johns wort

Dwarf St. Johns wort

Spotted St. Johns wort

ROCKROSE FAMILY

Narrow leaf pinweed

VIOLET FAMILY

Missouri violet

Birdsfoot violet

Smooth yellow violet

Field pansy

PASSION-FLOWER FAMILY

Maypop passion-flower

Yellow passion-flower

SOUR GUM FAMILY

Black tupelo

MELASTOMA FAMILY

Maryland meadow beauty

EVENING-PRIMROSE FAMILY

Tall gaura

Rattlebox

Evening-primrose

Goodman sundrops

Cut leaf evening-primrose

Showy primrose

WATER-MILFOIL FAMILY

Brazilian parrotfeather

UMBELLIFERAE

- **Chaerophyllum procumbens*
- **Chaerophyllum tainturieri*
- **Cicuta maculata*
- Eryngium yuccifolium*
- Polytaenia nuttallii*
- **Spermolepis divaricata*
- **Torilis arvensis*

CORNACEAE

- **Cornus drummondii*
- **Cornus florida*
- **Cornus obliqua*

ERICACEAE

- **Vaccinium arboreum*
- **Vaccinium stamineum*
- **Vaccinium vacillans*

PRIMULACEAE

- **Dodecatheon meadia*

SAPOTACEAE

- **Bumelia lanuginosa*

EBENACEAE

- **Diospyros virginiana*

OLEACEAE

- **Fraxinus americana* var.
americana
- Fraxinus pennsylvanica* var.
subintegerrima

GENTIANACEAE

- **Sabatia angularis*
- Sabatia campestris*

APOCYNACEAE

- **Amsonia hubrichtii*
- Amsonia tabernaemontana*
- **Apocynum cannabinum* var.
pubescens
- Trachelospermum difforme*

ASCLEPIADACEAE

- **Asclepias quadrifolia*
- **Asclepias tuberosa*
- **Asclepias verticillata*
- **Asclepias viridis*

CONVOLVULACEAE

- **Ipomoea pandurata*
- Ipomoea purpurea*

POLEMONIACEAE

- Gilia rubra*
- **Phlox pilosa* ssp.
ozarkana
- **Phlox pilosa* ssp.
pilosa

HYDROPHYLLACEAE

- Hydrolea ovata*
- Nemophila phacelioides*
- Phacelia glabra*
- **Phacelia hirsuta*

BORAGINACEAE

- **Heliotropium indicum*
- **Myosotis verna*

PARSLEY FAMILY

- Spreading chervil
- Tainturier chervil
- Water hemlock
- Button snakeroot
- Prairie parsley
- Rough fruit spermolepis
- Hedge parsley

DOGWOOD FAMILY

- Rough leaf dogwood
- Flowering dogwood
- Silky dogwood

HEATH FAMILY

- Tree huckleberry
- Deerberry
- Blueridge blueberry

PRIMROSE FAMILY

- Shootingstar

SAPODILLA FAMILY

- Chittamwood

EBONY FAMILY

- Persimmon

OLIVE FAMILY

- White ash

- Green ash

GENTIAN FAMILY

- Square stem rosegentian
- Prairie rosegentian

DOGBANE FAMILY

- Hubricht bluestar
- Low slimpod

- Hemp dogbane
- American starjasmine

MILKWEED FAMILY

- Four leaf milkweed
- Butterfly milkweed
- Whorled milkweed
- Green flower milkweed

CONVOLVULUS FAMILY

- Root morning-glory
- Morning-glory

PHLOX FAMILY

- Standing cypress

- Ozark downy phlox

- Downy phlox

WATERLEAF FAMILY

- Hairy hydrolea
- Large flower nemophila
- Smooth phacelia
- Hairy phacelia

BORAGE FAMILY

- Turnsole
- Early scorpiongrass

VERBENACEAE

- **Callicarpa americana*
- Verbena bracteata*
- **Verbena canadensis*
- **Verbena stricta*
- **Verbena urticifolia*

LABIATAE

- **Monarda russelliana*
- Monarda citriodora*
- **Monarda fistulosa*
- **Monarda virgata*
- **Prunella vulgaris*
- **Pycnanthemum tenuifolium*
- **Salvia azurea* var.
grandiflora
- **Salvia lyrata*
- **Scutellaria ovata*
- Scutellaria parvula* var.
australis

SOLANACEAE

- Physalis virginiana*
- Solanum americanum*
- **Solanum carolinense*
- Solanum elaeagnifolium*
- **Solanum rostratum*

SCROPHULARIACEAE

- **Agalinis fasciculata*
- **Agalinis heterophylla*
- **Agalinis tenuifolia*
- **Aureolaria grandiflora*
- **Bacopa rotundifolia*
- Buchnera americana*
- Castilleja coccinea*
- Castilleja indivisa*
- **Collinsia violacea*
- **Linaria canadensis* var.
texana
- Lindernia dubia*
- Mimulus alatus*
- **Pedicularis canadensis*
- Penstemon digitalis*
- Penstemon tubiflorus*
- **Verbascum blattaria* forma
albiflora
- **Verbascum thapsus*
- Veronica arvensis*

BIGNONIACEAE

- **Campsis radicans*
- **Catalpa speciosa*

ACANTHACEAE

- **Justicia americana*
- **Ruellia humilis*
- Ruellia strepens* forma
cleistantha

PLANTAGINACEAE

- **Plantago aristata*
- **Plantago virginica*

RUBIACEAE

- **Cephalanthus occidentalis*

VERVAIN FAMILY

- French mulberry
- Large bract vervain
- Rose vervain
- Woolly vervain
- White vervain

MINT FAMILY

- Russell beebalm
- Lemon beebalm
- Long flower horsemint
- Branched beebalm
- Selfheal
- Slender mountainmint

- Azure sage
- Lyre leaf sage
- Egg leaf skullcap

- Small skullcap

NIGHTSHADE FAMILY

- Virginia groundcherry
- Blue flower nightshade
- Horsenettle
- Silver leaf nightshade
- Buffalobur

FIGWORT FAMILY

- Fascicled agalinis
- Prairie agalinis
- Slender agalinis
- False foxglove
- Water hyssop
- American bluehearts
- Indian paintbrush
- Texas paintbrush
- Violet collinsia

- Old field toadflax
- Yellow seed falsepimpernel
- Sharp wing monkeyflower
- Common lousewort
- Beardtongue
- Tube penstemon

- Moth mullein
- Flannel mullein
- Common speedwell

BIGNONIA FAMILY

- Common trumpetvine
- Catalpa

ACANTHUS FAMILY

- American water-willow
- Wild petunia

- Limestone ruellia

PLANTAIN FAMILY

- Bracted plantain
- Paleseed plantain

MADDER FAMILY

- Buttonbush

Diodia teres
 **Diodia virginiana*
 **Galium pilosum* var.
 pilosum
Hedyotis crassifolia
Hedyotis nigricans
Sherardia arvensis
Spermacoce glabra
CAPRIFOLIACEAE
 **Lonicera flava*
 **Lonicera japonica*
 **Sambucus canadensis*
 **Symphoricarpos orbiculatus*
 **Viburnum rufidulum*
VALERIANACEAE
 **Valerianella nuttallii*
 **Valerianella radiata*
CAMPANULACEAE
 **Lobelia cardinalis*
 **Lobelia puberula*
Triodanis lamprosperma
Triodanis perfoliata
COMPOSITAE
 **Achillea millefolium* var.
 lanulosa
 **Ambrosia bidentata*
 **Ambrosia psilostachya*
 **Antennaria plantaginifolia*
Anthemis cotula
Aster azureus
 **Aster ericoides* var.
 ericoides
 **Aster lateriflorus* var.
 flagellaris
 **Aster paludosus* var.
 hemisphericus
 **Aster patens* var.
 patentissimus
 **Aster sagittifolius*
 **Aster turbinellus*
 **Bidens polylepis*
 **Boltonia diffusa*
Cacalia tuberosa
Chaetopappa asteroides
 **Chrysopsis pilosa*
Cirsium altissimum
 **Cirsium carolinianum*
Cirsium iowense
 **Conyza canadensis*
 **Coreopsis grandiflora* var.
 harveyana
 **Coreopsis tinctoria*
Coreopsis tripteris
Crepis pulchra
Dracopsis amplexicaulis
Echinacea pallida
 **Elephantopus carolinianus*
 **Erigeron strigosus*
Erigeron tenuis
 **Eupatorium coelestinum*
 **Eupatorium rugosum*

Rough buttonweed
 Large buttonweed

 Hairy bedstraw
 Small bluets
 Prairie bluets
 Field madder
 Smooth buttonplant
HONEYSUCKLE FAMILY
 Yellow honeysuckle
 Japanese honeysuckle
 American elder
 Coralberry
 Southern blackhaw
VALERIAN FAMILY
 Nuttall cornsalad
 Beaked cornsalad
BLUEBELL FAMILY
 Cardinal flower
 Purple dewdrop
 Venus lookingglass
 Clasping Venus lookingglass
SUNFLOWER FAMILY

 Yarrow
 Lance leaf ragweed
 Western ragweed
 Plantain leaf pussytoes
 Mayweed camomile
 Sky-blue aster

 Heath aster

 Calico aster

 Single stem bog-aster

 Skydrop aster
 Arrow aster
 Prairie aster
 Awnless beggarticks
 Small head boltonia
 Indian plantain
 Leastdaisy
 Soft gold aster
 Tall thistle
 Soft thistle
 Iowa thistle
 Canada fleabane

 Cord leaf coreopsis
 Plains coreopsis
 Atlantic coreopsis
 Showy hawksbeard
 Clasping coneflower
 Pale echinacea
 Leafy elephantfoot
 Prairie fleabane
 Slender fleabane
 Mistflower
 White snakeroot

| | |
|--|--------------------------|
| * <i>Eupatorium serotinum</i> | Late flower thoroughwort |
| * <i>Euthamia gymnospermoides</i> | Bushy goldenrod |
| * <i>Gnaphalium obtusifolium</i> | Fragrant cudweed |
| * <i>Gnaphalium purpureum</i> | Purple cudweed |
| <i>Grindelia lanceolata</i> | Gumweed |
| * <i>Gutierrezia dracunculoides</i> | Broomweed |
| * <i>Helenium amarum</i> | Bitterweed |
| <i>Helenium flexuosum</i> | Purple head sneezeweed |
| * <i>Helianthus divaricatus</i> | Rough sunflower |
| * <i>Helianthus hirsutus</i> | Hairy sunflower |
| <i>Helianthus mollis</i> | Ashy sunflower |
| * <i>Heliopsis helianthoides</i> var.
<i>scabra</i> | Rough heliopsis |
| <i>Hieracium gronovii</i> | Hairy hawkweed |
| * <i>Krigia virginica</i> | Carolina dwarf dandelion |
| * <i>Liatris asper</i> var.
<i>intermedia</i> | Rough gayfeather |
| * <i>Liatris elegans</i> | Pink scale gayfeather |
| * <i>Liatris pycnostachya</i> | Hairy button snakeroot |
| * <i>Liatris squarrosa</i> var.
<i>hirsuta</i> | Scaly blazing star |
| * <i>Pyrrhopappus carolinianus</i> | Carolina false dandelion |
| * <i>Rudbeckia grandiflora</i> | Rough coneflower |
| * <i>Rudbeckia hirta</i> var.
<i>pulcherrima</i> | Black eyed Susan |
| * <i>Rudbeckia subtomentosa</i> | Sweet coneflower |
| * <i>Rudbeckia triloba</i> | Brown eyed Susan |
| <i>Senecio glabellus</i> | Butterweed |
| <i>Senecio obovatus</i> | Golden groundsel |
| * <i>Serinia oppositifolia</i> | Weedy dwarf dandelion |
| <i>Silphium asteriscus</i> | Starry rosinweed |
| * <i>Solidago canadensis</i> var.
<i>scabra</i> | Canada goldenrod |
| <i>Solidago gigantea</i> var.
<i>serotina</i> | Giant goldenrod |
| <i>Solidago missouriensis</i> | Missouri goldenrod |
| * <i>Solidago petiolaris</i> | Downy goldenrod |
| * <i>Solidago radula</i> | Rough goldenrod |
| * <i>Verbesina virginica</i> | White crownbeard |
| * <i>Vernonia baldwinii</i> | Baldwin ironweed |
| * <i>Vernonia lettermanii</i> | Letterman ironweed |

References

- Curtis, N. M., Jr., and Ham, W. E., 1972, Geomorphic provinces of Oklahoma, *in* Johnson, K. S., Branson, C. C., Curtis, N. M., Jr., Ham, W. E., Harrison, W. E., Marcher, M. V., and Roberts, J. F., *Geology and earth resources of Oklahoma—An atlas of maps and cross sections: Oklahoma Geological Survey Educational Publication 1*, p. 3.
- Johnson, K. S., Branson, C. C., Curtis, N. M., Jr., Ham, W. E., Marcher, M. V., and Roberts, J. F., 1972, *Geology and earth resources of Oklahoma—An atlas of maps and cross sections: Oklahoma Geological Survey Educational Publication 1*, 8 p.
- Raisz, E. J., 1957, *Map of the landforms of the United States* [6th revised edition; originally published in Atwood, W. W., 1940, *Physiographic provinces of North America*].
- Russell, D. T., 1958, *Guidebook to Robbers Cave State Park and Camp Tom Hale, Latimer County, Oklahoma: Oklahoma Geological Survey Guidebook 7*, 23 p.

Glossary of Geological Terms

anticline—An upfold of strata the center of which is the "axis." The strata dip away from the axis like the roof of a house. The core of the fold contains stratigraphically older rocks. The opposite of a syncline.

bedding plane—A characteristic of sedimentary rocks. Bedding planes are subparallel planes along which individual beds may separate.

clastic rock—Sedimentary rock consisting of fragments derived from preexisting rocks that were broken down and then transported to another place of deposition.

columnar section—A drawing or graphic representation showing in a vertical column the sequence of rock units found in a specified area.

cuesta—A ridge with one slope long and gentle and the other slope steep. *Cuesta* is a Spanish word meaning flank or slope of a hill. Cuestas are caused by dipping rock strata.

dip—The angle at which a rock stratum or fault surface is inclined from the horizontal. Dip is always at right angles (90°) to the strike of a stratum.

epoch—Subdivision of a period of time. Rocks formed during an epoch constitute a series.

era—Greatest of all geologic time divisions. Consists of two or more geologic periods.

erosion—The natural process of weathering, disintegration, dissolution, and removal of rock and earth material, mainly by water and wind.

fault—A fracture or a zone of fractures in rock with displacement of the sides relative to one another either horizontally, vertically, or both.

flood plain—A relatively flat area close to a stream. It is made of sediment carried over the stream banks during times of flood.

fold—A curve or bend in rock strata, usually as a result of deformation.

formation—The fundamental stratigraphic unit used for geologic mapping. It may include several rock types. Formations sometimes are divided into members and beds.

fossil—The remains or traces of an animal or plant that have been preserved by natural causes in the Earth's crustal rocks. The term does not include an organism that has been buried during historic time.

geologic time—History of the Earth divided into eras, periods, and epochs.

group—Two or more formations with similar characteristics or other affinities.

hogback—The outcropping edge of a cuesta. Its slopes are roughly equal.

igneous rock—Rock formed by cooling and solidification of hot molten material called magma. Magma that flows onto the surface of the earth (lava) cools rapidly to form fine-grained rocks, whereas magma that solidifies several miles beneath the surface cools slowly to form coarser grained rocks.

joint—A fracture in rock along which no displacement has occurred.

marine—Of or belonging to the sea.

member—*see* formation

metamorphic rock—Rock that has been changed through intense heat, high pressure, or contact with chemically active fluids from magma. Slate and phyllite are examples of metamorphosed shale, quartzite metamorphosed sandstone, and marble metamorphosed limestone.

nonclastic rock—A sedimentary rock whose deposition was caused by chemical or biological action.

organic deposits—Deposits formed from the remains of living matter such as plants and animals.

orogeny—A movement or movements of the Earth's crust (folding and faulting) that cause mountains to form.

paleobotany—The science of fossil plants.

paleontology—The science of fossils, both plant and animal.

paleozoology—The science of fossil animals. Its two divisions are vertebrate and invertebrate paleozoology.

period—Subdivision of an era. Rocks formed during a period constitute a system.

sediment—Bits of rock, often the result of weathering, that accumulate in layers. Sand, gravel, silt, and mud are some examples of sediment, which can be transported or deposited by air, water, or ice.

sedimentary rock—Rock formed by the compaction and cementing of sediments deposited in water or from air. Sediments may consist of rock or mineral fragments of various sizes (mud, sand, gravel), the remains of animals or plants, the products of chemical action or evaporation, or mixtures of these materials. Sedimentary rocks typically have a layered structure known as bedding or stratification.

series—*see* epoch

slump—A mass of rock or unconsolidated material of any size that has slipped downward by gravity from its original position.

stratum—A single sedimentary bed or layer. The plural form is strata.

strike—The direction or trend of a rock stratum or fault surface as it intersects the horizontal.

structural geology—Study of the structural features of rocks (folds and faults) and the causes of their deformation.

syncline—A fold in rock layers in which the strata dip inward from both sides toward the axis. A downfold. The core of the fold contains stratigraphically younger rocks. The opposite of an anticline.


system—*see* period

terrestrial—Consisting of or pertaining to land as distinct from water. Examples of terrestrial deposits are flood plains of streams and rivers. These clastic sediments are deposited by flood waters.

topography—The physical features of a region; the shape of the land's surface.

water table—The upper limit of the portion of the ground that is wholly saturated with water.

weathering—The group of processes—such as the chemical action of water and air and of plants and bacteria, and the mechanical action of changes of temperature—whereby rocks exposed to the weather change in character, decay, and finally crumble into soil.



By
Dearl T. Russell and Arthur J. Myers
1985

